

Light-induced Gauge Potentials for Cold Atoms

Julius Ruseckas Gediminas Juzeliūnas

Institute of Theoretical Physics and Astronomy, Vilnius University, Lithuania

August 30, 2010

- 1 Motivation
- 2 Some aspects of adiabatic approximation
- 3 Abelian effective potentials
- 4 Non-Abelian effective potentials for tripod coupling scheme
 - Rashba-type Hamiltonian with spin $1/2$
- 5 Non-Abelian fields in N -pod schemes
 - Rashba-type Hamiltonian with spin 1

- 1 Motivation
- 2 Some aspects of adiabatic approximation
- 3 Abelian effective potentials
- 4 Non-Abelian effective potentials for tripod coupling scheme
 - Rashba-type Hamiltonian with spin $1/2$
- 5 Non-Abelian fields in N -pod schemes
 - Rashba-type Hamiltonian with spin 1

- 1 Motivation
- 2 Some aspects of adiabatic approximation
- 3 Abelian effective potentials
- 4 Non-Abelian effective potentials for tripod coupling scheme
 - Rashba-type Hamiltonian with spin $1/2$
- 5 Non-Abelian fields in N -pod schemes
 - Rashba-type Hamiltonian with spin 1

- 1 Motivation
- 2 Some aspects of adiabatic approximation
- 3 Abelian effective potentials
- 4 Non-Abelian effective potentials for tripod coupling scheme
 - Rashba-type Hamiltonian with spin $1/2$
- 5 Non-Abelian fields in N -pod schemes
 - Rashba-type Hamiltonian with spin 1

- 1 Motivation
- 2 Some aspects of adiabatic approximation
- 3 Abelian effective potentials
- 4 Non-Abelian effective potentials for tripod coupling scheme
 - Rashba-type Hamiltonian with spin $1/2$
- 5 Non-Abelian fields in N -pod schemes
 - Rashba-type Hamiltonian with spin 1

Why effective magnetic field for atoms?

Atomic physics \iff Solid state physics:

- Degenerate Fermi gas \iff Electrons in solids
- Atoms in optical lattices

Advantages and disadvantages of cold atoms

- **Advantage:** Freedom in changing experimental parameters that are often inaccessible in standard solid state physics
- **Disadvantage:** Trapped atoms are electrically neutral particles. Direct analogy with magnetic properties of solids is not necessarily straightforward

Why effective magnetic field for atoms?

Atomic physics \iff Solid state physics:

- Degenerate Fermi gas \iff Electrons in solids
- Atoms in optical lattices

Advantages and disadvantages of cold atoms

- **Advantage:** Freedom in changing experimental parameters that are often inaccessible in standard solid state physics
- **Disadvantage:** Trapped atoms are electrically neutral particles. Direct analogy with magnetic properties of solids is not necessarily straightforward

Why effective magnetic field for atoms?

Atomic physics \iff Solid state physics:

- Degenerate Fermi gas \iff Electrons in solids
- Atoms in optical lattices

Advantages and disadvantages of cold atoms

- **Advantage:** Freedom in changing experimental parameters that are often inaccessible in standard solid state physics
- **Disadvantage:** Trapped atoms are electrically neutral particles. Direct analogy with magnetic properties of solids is not necessarily straightforward

Why effective magnetic field for atoms?

Atomic physics \iff Solid state physics:

- Degenerate Fermi gas \iff Electrons in solids
- Atoms in optical lattices

Advantages and disadvantages of cold atoms

- **Advantage:** Freedom in changing experimental parameters that are often inaccessible in standard solid state physics
- **Disadvantage:** Trapped atoms are electrically neutral particles. Direct analogy with magnetic properties of solids is not necessarily straightforward

Why effective magnetic field for atoms?

Atomic physics \iff Solid state physics:

- Degenerate Fermi gas \iff Electrons in solids
- Atoms in optical lattices

Advantages and disadvantages of cold atoms

- **Advantage:** Freedom in changing experimental parameters that are often inaccessible in standard solid state physics
- **Disadvantage:** Trapped atoms are electrically neutral particles. Direct analogy with magnetic properties of solids is not necessarily straightforward

Analogies with the elementary particle physics

Cold atomic gases are an analog not only to the solid state physics. Creation of the effective gauge potentials allows for the motion of cold atoms to be described by equations that usually appear in the elementary particle physics.

- Non-Abelian gauge potentials
- Magnetic monopole
- Ultrarelativistic Dirac fermions
- Zitterbewegung
- Negative reflection

Analogies with the elementary particle physics

Cold atomic gases are an analog not only to the solid state physics. Creation of the effective gauge potentials allows for the motion of cold atoms to be described by equations that usually appear in the elementary particle physics.

- Non-Abelian gauge potentials
- Magnetic monopole
- Ultrarelativistic Dirac fermions
- Zitterbewegung
- Negative reflection

Analogies with the elementary particle physics

Cold atomic gases are an analog not only to the solid state physics. Creation of the effective gauge potentials allows for the motion of cold atoms to be described by equations that usually appear in the elementary particle physics.

- Non-Abelian gauge potentials
- Magnetic monopole
- Ultrarelativistic Dirac fermions
- Zitterbewegung
- Negative reflection

Analogies with the elementary particle physics

Cold atomic gases are an analog not only to the solid state physics. Creation of the effective gauge potentials allows for the motion of cold atoms to be described by equations that usually appear in the elementary particle physics.

- Non-Abelian gauge potentials
- Magnetic monopole
- Ultrarelativistic Dirac fermions
- Zitterbewegung
- Negative reflection

Analogies with the elementary particle physics

Cold atomic gases are an analog not only to the solid state physics. Creation of the effective gauge potentials allows for the motion of cold atoms to be described by equations that usually appear in the elementary particle physics.

- Non-Abelian gauge potentials
- Magnetic monopole
- Ultrarelativistic Dirac fermions
- Zitterbewegung
- Negative reflection

Analogies with the elementary particle physics

Cold atomic gases are an analog not only to the solid state physics. Creation of the effective gauge potentials allows for the motion of cold atoms to be described by equations that usually appear in the elementary particle physics.

- Non-Abelian gauge potentials
- Magnetic monopole
- Ultrarelativistic Dirac fermions
- Zitterbewegung
- Negative reflection

Magnetic field and rotation

- Coriolis force:

$$\mathbf{F}_C = 2m\mathbf{v} \times \boldsymbol{\Omega}$$

- Lorentz force:

$$\mathbf{F}_L = q\mathbf{v} \times \mathbf{B}$$

Rotation is similar to the magnetic field.

- Coriolis force:

$$\mathbf{F}_C = 2m\mathbf{v} \times \boldsymbol{\Omega}$$

- Lorentz force:

$$\mathbf{F}_L = q\mathbf{v} \times \mathbf{B}$$

Rotation is similar to the magnetic field.

Ways to create effective magnetic field for cold atoms

- **Rotation** — usual method to create effective magnetic field
 - Constant effective magnetic field $B_{\text{eff}} \sim \Omega$
 - Trapping frequency $\omega_{\text{eff}} = \omega - \Omega$
 - Effective magnetic field acts on atoms in the same way
- **Optical lattices** having asymmetry in the atomic transitions between the lattice sites.
 - **Abelian** effective gauge potentials
 - D. Jaksch and P. Zoller, New J. Phys. **5**, 56 (2003)
 - E. Mueller, Phys. Rev. A **70**, 04163(R) (2004)
 - A. S. Sørensen, E. Demler, and M. D. Lukin, Phys. Rev. Lett. **94**, 086803 (2005)
 - **Non-Abelian** effective gauge potentials in optical lattices
 - K. Osterloh, M. Baig, L. Santos, P. Zoller, and M. Lewenstein, Phys. Rev. Lett. **95**, 010403 (2005)

Ways to create effective magnetic field for cold atoms

- **Rotation** — usual method to create effective magnetic field
 - Constant effective magnetic field $B_{\text{eff}} \sim \Omega$
 - Trapping frequency $\omega_{\text{eff}} = \omega - \Omega$
 - Effective magnetic field acts on atoms in the same way
- **Optical lattices** having asymmetry in the atomic transitions between the lattice sites.
 - **Abelian** effective gauge potentials
 - D. Jaksch and P. Zoller, New J. Phys. **5**, 56 (2003)
 - E. Mueller, Phys. Rev. A **70**, 04163(R) (2004)
 - A. S. Sørensen, E. Demler, and M. D. Lukin, Phys. Rev. Lett. **94**, 086803 (2005)
 - **Non-Abelian** effective gauge potentials in optical lattices
 - K. Osterloh, M. Baig, L. Santos, P. Zoller, and M. Lewenstein, Phys. Rev. Lett. **95**, 010403 (2005)

Ways to create effective magnetic field for cold atoms

- **Rotation** — usual method to create effective magnetic field
 - Constant effective magnetic field $B_{\text{eff}} \sim \Omega$
 - Trapping frequency $\omega_{\text{eff}} = \omega - \Omega$
 - Effective magnetic field acts on atoms in the same way
- **Optical lattices** having asymmetry in the atomic transitions between the lattice sites.
 - **Abelian** effective gauge potentials
 - D. Jaksch and P. Zoller, New J. Phys. **5**, 56 (2003)
 - E. Mueller, Phys. Rev. A **70**, 04163(R) (2004)
 - A. S. Sørensen, E. Demler, and M. D. Lukin, Phys. Rev. Lett. **94**, 086803 (2005)
 - **Non-Abelian** effective gauge potentials in optical lattices
 - K. Osterloh, M. Baig, L. Santos, P. Zoller, and M. Lewenstein, Phys. Rev. Lett. **95**, 010403 (2005)

Ways to create effective magnetic field for cold atoms

- **Rotation** — usual method to create effective magnetic field
 - Constant effective magnetic field $B_{\text{eff}} \sim \Omega$
 - Trapping frequency $\omega_{\text{eff}} = \omega - \Omega$
 - Effective magnetic field acts on atoms in the same way
- **Optical lattices** having asymmetry in the atomic transitions between the lattice sites.
 - **Abelian** effective gauge potentials
 - D. Jaksch and P. Zoller, New J. Phys. **5**, 56 (2003)
 - E. Mueller, Phys. Rev. A **70**, 04163(R) (2004)
 - A. S. Sørensen, E. Demler, and M. D. Lukin, Phys. Rev. Lett. **94**, 086803 (2005)
 - **Non-Abelian** effective gauge potentials in optical lattices
 - K. Osterloh, M. Baig, L. Santos, P. Zoller, and M. Lewenstein, Phys. Rev. Lett. **95**, 010403 (2005)

Ways to create effective magnetic field for cold atoms

- **Rotation** — usual method to create effective magnetic field
 - Constant effective magnetic field $B_{\text{eff}} \sim \Omega$
 - Trapping frequency $\omega_{\text{eff}} = \omega - \Omega$
 - Effective magnetic field acts on atoms in the same way
- **Optical lattices** having asymmetry in the atomic transitions between the lattice sites.
 - **Abelian** effective gauge potentials
 - D. Jaksch and P. Zoller, New J. Phys. **5**, 56 (2003)
 - E. Mueller, Phys. Rev. A **70**, 04163(R) (2004)
 - A. S. Sørensen, E. Demler, and M. D. Lukin, Phys. Rev. Lett. **94**, 086803 (2005)
 - **Non-Abelian** effective gauge potentials in optical lattices
 - K. Osterloh, M. Baig, L. Santos, P. Zoller, and M. Lewenstein, Phys. Rev. Lett. **95**, 010403 (2005)

Ways to create effective magnetic field for cold atoms

- **Rotation** — usual method to create effective magnetic field
 - Constant effective magnetic field $B_{\text{eff}} \sim \Omega$
 - Trapping frequency $\omega_{\text{eff}} = \omega - \Omega$
 - Effective magnetic field acts on atoms in the same way
- **Optical lattices** having asymmetry in the atomic transitions between the lattice sites.
 - **Abelian** effective gauge potentials
 - D. Jaksch and P. Zoller, New J. Phys. **5**, 56 (2003)
 - E. Mueller, Phys. Rev. A **70**, 04163(R) (2004)
 - A. S. Sørensen, E. Demler, and M. D. Lukin, Phys. Rev. Lett. **94**, 086803 (2005)
 - **Non-Abelian** effective gauge potentials in optical lattices
 - K. Osterloh, M. Baig, L. Santos, P. Zoller, and M. Lewenstein, Phys. Rev. Lett. **95**, 010403 (2005)

Ways to create effective magnetic field for cold atoms

- **Rotation** — usual method to create effective magnetic field
 - Constant effective magnetic field $B_{\text{eff}} \sim \Omega$
 - Trapping frequency $\omega_{\text{eff}} = \omega - \Omega$
 - Effective magnetic field acts on atoms in the same way
- **Optical lattices** having asymmetry in the atomic transitions between the lattice sites.
 - **Abelian** effective gauge potentials
 - D. Jaksch and P. Zoller, New J. Phys. **5**, 56 (2003)
 - E. Mueller, Phys. Rev. A **70**, 04163(R) (2004)
 - A. S. Sørensen, E. Demler, and M. D. Lukin, Phys. Rev. Lett. **94**, 086803 (2005)
 - **Non-Abelian** effective gauge potentials in optical lattices
 - K. Osterloh, M. Baig, L. Santos, P. Zoller, and M. Lewenstein, Phys. Rev. Lett. **95**, 010403 (2005)

Effective magnetic field induced by EIT

Effective gauge potentials can be created using light beams with non-zero relative orbital angular momentum (OAM) in the **EIT** configuration.

Advantages

- No rotation
- No need for optical lattice

Abelian gauge fields:

- G. Juzeliūnas and P. Öhberg, Phys. Rev. Lett. **93**, 033602 (2004).

Non-Abelian gauge fields:

- J. Ruseckas, G. Juzeliūnas, P. Öhberg, and M. Fleischhauer Phys. Rev. Lett. **95**, 010404 (2005)

Effective magnetic field induced by EIT

Effective gauge potentials can be created using light beams with non-zero relative orbital angular momentum (OAM) in the **EIT** configuration.

Advantages

- No rotation
- No need for optical lattice

Abelian gauge fields:

- G. Juzeliūnas and P. Öhberg, Phys. Rev. Lett. **93**, 033602 (2004).

Non-Abelian gauge fields:

- J. Ruseckas, G. Juzeliūnas, P. Öhberg, and M. Fleischhauer Phys. Rev. Lett. **95**, 010404 (2005)

Effective magnetic field induced by EIT

Effective gauge potentials can be created using light beams with non-zero relative orbital angular momentum (OAM) in the **EIT** configuration.

Advantages

- No rotation
- No need for optical lattice

Abelian gauge fields:

- G. Juzeliūnas and P. Öhberg, Phys. Rev. Lett. **93**, 033602 (2004).

Non-Abelian gauge fields:

- J. Ruseckas, G. Juzeliūnas, P. Öhberg, and M. Fleischhauer Phys. Rev. Lett. **95**, 010404 (2005)

Effective gauge potentials can be created using light beams with non-zero relative orbital angular momentum (OAM) in the **EIT** configuration.

Advantages

- No rotation
- No need for optical lattice

Abelian gauge fields:

- G. Juzeliūnas and P. Öhberg, Phys. Rev. Lett. **93**, 033602 (2004).

Non-Abelian gauge fields:

- J. Ruseckas, G. Juzeliūnas, P. Öhberg, and M. Fleischhauer Phys. Rev. Lett. **95**, 010404 (2005)

Adiabatic Approximation

- The full atomic Hamiltonian

$$\hat{H} = \frac{\hat{p}^2}{2M} + \hat{V}(\mathbf{r}) + \hat{H}_0(\mathbf{r}, t).$$

- $\hat{H}_0(\mathbf{r}, t)$ — the Hamiltonian for the electronic (fast) degrees of freedom,
- $\hat{p}^2/2M + \hat{V}(\mathbf{r})$ — the Hamiltonian for center of mass (slow) degrees of freedom.
- $\hat{V}(\mathbf{r})$ — the external trapping potential.
- $\hat{H}_0(\mathbf{r}, t)$ has eigenfunctions $|\chi_n(\mathbf{r}, t)\rangle$ with eigenvalues $\varepsilon(\mathbf{r}, t)$.
- Full atomic wave function

$$|\Phi\rangle = \sum_n \Psi_n(\mathbf{r}, t) |\chi_n(\mathbf{r}, t)\rangle.$$

Adiabatic Approximation

- The full atomic Hamiltonian

$$\hat{H} = \frac{\hat{p}^2}{2M} + \hat{V}(\mathbf{r}) + \hat{H}_0(\mathbf{r}, t).$$

- $\hat{H}_0(\mathbf{r}, t)$ — the Hamiltonian for the electronic (**fast**) degrees of freedom,
- $\hat{p}^2/2M + \hat{V}(\mathbf{r})$ — the Hamiltonian for center of mass (**slow**) degrees of freedom.
- $\hat{V}(\mathbf{r})$ — the external trapping potential.
- $\hat{H}_0(\mathbf{r}, t)$ has eigenfunctions $|\chi_n(\mathbf{r}, t)\rangle$ with eigenvalues $\varepsilon(\mathbf{r}, t)$.
- Full atomic wave function

$$|\Phi\rangle = \sum_n \Psi_n(\mathbf{r}, t) |\chi_n(\mathbf{r}, t)\rangle.$$

Adiabatic Approximation

- The full atomic Hamiltonian

$$\hat{H} = \frac{\hat{p}^2}{2M} + \hat{V}(\mathbf{r}) + \hat{H}_0(\mathbf{r}, t).$$

- $\hat{H}_0(\mathbf{r}, t)$ — the Hamiltonian for the electronic (**fast**) degrees of freedom,
- $\hat{p}^2/2M + \hat{V}(\mathbf{r})$ — the Hamiltonian for center of mass (**slow**) degrees of freedom.
- $\hat{V}(\mathbf{r})$ — the external trapping potential.
- $\hat{H}_0(\mathbf{r}, t)$ has eigenfunctions $|\chi_n(\mathbf{r}, t)\rangle$ with eigenvalues $\varepsilon(\mathbf{r}, t)$.
- Full atomic wave function

$$|\Phi\rangle = \sum_n \Psi_n(\mathbf{r}, t) |\chi_n(\mathbf{r}, t)\rangle.$$

Adiabatic Approximation

- The full atomic Hamiltonian

$$\hat{H} = \frac{\hat{p}^2}{2M} + \hat{V}(\mathbf{r}) + \hat{H}_0(\mathbf{r}, t).$$

- $\hat{H}_0(\mathbf{r}, t)$ — the Hamiltonian for the electronic (**fast**) degrees of freedom,
- $\hat{p}^2/2M + \hat{V}(\mathbf{r})$ — the Hamiltonian for center of mass (**slow**) degrees of freedom.
- $\hat{V}(\mathbf{r})$ — the external trapping potential.
- $\hat{H}_0(\mathbf{r}, t)$ has eigenfunctions $|\chi_n(\mathbf{r}, t)\rangle$ with eigenvalues $\varepsilon(\mathbf{r}, t)$.
- Full atomic wave function

$$|\Phi\rangle = \sum_n \Psi_n(\mathbf{r}, t) |\chi_n(\mathbf{r}, t)\rangle.$$

Adiabatic Approximation

- The full atomic Hamiltonian

$$\hat{H} = \frac{\hat{p}^2}{2M} + \hat{V}(\mathbf{r}) + \hat{H}_0(\mathbf{r}, t).$$

- $\hat{H}_0(\mathbf{r}, t)$ — the Hamiltonian for the electronic (**fast**) degrees of freedom,
- $\hat{p}^2/2M + \hat{V}(\mathbf{r})$ — the Hamiltonian for center of mass (**slow**) degrees of freedom.
- $\hat{V}(\mathbf{r})$ — the external trapping potential.
- $\hat{H}_0(\mathbf{r}, t)$ has eigenfunctions $|\chi_n(\mathbf{r}, t)\rangle$ with eigenvalues $\varepsilon(\mathbf{r}, t)$.
- Full atomic wave function

$$|\Phi\rangle = \sum_n \Psi_n(\mathbf{r}, t) |\chi_n(\mathbf{r}, t)\rangle.$$

Adiabatic Approximation

- The full atomic Hamiltonian

$$\hat{H} = \frac{\hat{p}^2}{2M} + \hat{V}(\mathbf{r}) + \hat{H}_0(\mathbf{r}, t).$$

- $\hat{H}_0(\mathbf{r}, t)$ — the Hamiltonian for the electronic (**fast**) degrees of freedom,
- $\hat{p}^2/2M + \hat{V}(\mathbf{r})$ — the Hamiltonian for center of mass (**slow**) degrees of freedom.
- $\hat{V}(\mathbf{r})$ — the external trapping potential.
- $\hat{H}_0(\mathbf{r}, t)$ has eigenfunctions $|\chi_n(\mathbf{r}, t)\rangle$ with eigenvalues $\varepsilon(\mathbf{r}, t)$.
- Full atomic wave function

$$|\Phi\rangle = \sum_n \Psi_n(\mathbf{r}, t) |\chi_n(\mathbf{r}, t)\rangle.$$

Adiabatic Approximation

Substituting into the Schrödinger equation $i\hbar\partial/\partial t|\Phi\rangle = \hat{H}|\Phi\rangle$ one can write the equation for the coefficients $\Psi_n(\mathbf{r}, t)$ in the form

$$i\hbar\frac{\partial}{\partial t}\Psi = \left[\frac{1}{2M}(-i\hbar\nabla - \mathbf{A})^2 + V + \beta \right] \Psi,$$

where

$$\Psi = \begin{pmatrix} \Psi_1 \\ \dots \\ \Psi_n \end{pmatrix},$$

$$\mathbf{A}_{n,n'} = i\hbar\langle\chi_n(\mathbf{r}, t)|\nabla\chi_{n'}(\mathbf{r}, t)\rangle,$$

$$V_{n,n'} = \varepsilon(\mathbf{r}, t)\delta_{n,n'} + \langle\chi_n(\mathbf{r}, t)|\hat{V}(\mathbf{r})|\chi_{n'}(\mathbf{r}, t)\rangle,$$

$$\beta_{n,n'} = -i\hbar\langle\chi_n(\mathbf{r}, t)|\frac{\partial}{\partial t}\chi_{n'}(\mathbf{r}, t)\rangle.$$

Non-degenerate states

The first state is well separated from the rest. Off-diagonal terms are neglected.

$$i\hbar \frac{\partial}{\partial t} \Psi_1 = \left[\frac{1}{2M} (-i\hbar \nabla - \mathbf{A})^2 + V + \phi + \beta \right] \Psi_1,$$

where

$$\mathbf{A} = \mathbf{A}_{1,1},$$

$$V = V_{1,1},$$

$$\phi = \frac{1}{2M} \sum_{n \neq 1} \mathbf{A}_{1,n} \cdot \mathbf{A}_{n,1}.$$

Adiabatic Approximation

Degenerate states

The first q dressed states are degenerate and these levels are well separated from the remaining $N - q$

$$i\hbar \frac{\partial}{\partial t} \tilde{\Psi} = \left[\frac{1}{2M} (-i\hbar \nabla - \mathbf{A})^2 + V + \phi + \beta \right] \tilde{\Psi},$$

where \mathbf{A} and ϕ are truncated $q \times q$ matrices,

$$\phi_{n,n'} = \frac{1}{2M} \sum_{m=q+1}^N \mathbf{A}_{n,m} \cdot \mathbf{A}_{m,n'}.$$

The effective vector potential \mathbf{A} is called the **Mead-Berry connection**.
The effective scalar potential ϕ is called the **Born-Huang potential**.

Non-degenerate states

We have freedom of choosing the phase of the adiabatic states

$$|\chi_n(\mathbf{r}, t)\rangle \rightarrow e^{-\frac{i}{\hbar}u_n(\mathbf{r}, t)}|\chi_n(\mathbf{r}, t)\rangle.$$

The transformation of the potentials

$$\mathbf{A} \rightarrow \mathbf{A} + \nabla u_1,$$

$$\phi \rightarrow \phi - \frac{\partial}{\partial t} u_1.$$

Non-degenerate states

We have freedom of choosing the phase of the adiabatic states

$$|\chi_n(\mathbf{r}, t)\rangle \rightarrow e^{-\frac{i}{\hbar}u_n(\mathbf{r}, t)}|\chi_n(\mathbf{r}, t)\rangle.$$

The transformation of the potentials

$$\mathbf{A} \rightarrow \mathbf{A} + \nabla u_1,$$

$$\phi \rightarrow \phi - \frac{\partial}{\partial t}u_1.$$

Degenerate states

The adiabatic basis can be changed by a local unitary transformation $U(\mathbf{r}, t)$

$$\tilde{\Psi} \rightarrow U(\mathbf{r}, t)\Psi.$$

The transformation of the potentials

$$\mathbf{A} \rightarrow U\mathbf{A}U^\dagger - i\hbar(\nabla U)U^\dagger,$$

$$\phi \rightarrow U\phi U^\dagger + i\hbar\frac{\partial U}{\partial t}U^\dagger.$$

The Berry connection \mathbf{A} is related to a **curvature** \mathbf{B} as

$$B_i = \frac{1}{2}\epsilon_{ikl}F_{kl}, \quad F_{kl} = \partial_k A_l - \partial_l A_k - \frac{i}{\hbar}[A_k, A_l].$$

Degenerate states

The adiabatic basis can be changed by a local unitary transformation $U(\mathbf{r}, t)$

$$\tilde{\Psi} \rightarrow U(\mathbf{r}, t)\Psi.$$

The transformation of the potentials

$$\mathbf{A} \rightarrow U\mathbf{A}U^\dagger - i\hbar(\nabla U)U^\dagger,$$

$$\phi \rightarrow U\phi U^\dagger + i\hbar\frac{\partial U}{\partial t}U^\dagger.$$

The Berry connection \mathbf{A} is related to a **curvature** \mathbf{B} as

$$B_i = \frac{1}{2}\epsilon_{ikl}F_{kl}, \quad F_{kl} = \partial_k A_l - \partial_l A_k - \frac{i}{\hbar}[A_k, A_l].$$

Degenerate states

The adiabatic basis can be changed by a local unitary transformation $U(\mathbf{r}, t)$

$$\tilde{\Psi} \rightarrow U(\mathbf{r}, t)\tilde{\Psi}.$$

The transformation of the potentials


$$\mathbf{A} \rightarrow U\mathbf{A}U^\dagger - i\hbar(\nabla U)U^\dagger,$$

$$\phi \rightarrow U\phi U^\dagger + i\hbar\frac{\partial U}{\partial t}U^\dagger.$$

The Berry connection \mathbf{A} is related to a **curvature** \mathbf{B} as

$$B_i = \frac{1}{2}\epsilon_{ikl}F_{kl}, \quad F_{kl} = \partial_k A_l - \partial_l A_k - \frac{i}{\hbar}[A_k, A_l].$$

Λ -type Atoms




Probe beam: $\Omega_p = \mu_{13}E_p$
Control beam: $\Omega_c = \mu_{23}E_c$

Dark state

$$|D\rangle \sim \Omega_c|1\rangle - \Omega_p|2\rangle$$

Destructive interference,
cancellation of absorption
— EIT

Λ -type Atoms



Probe beam: $\Omega_p = \mu_{13} E_p$
Control beam: $\Omega_c = \mu_{23} E_c$

Dark state

$$|D\rangle \sim \Omega_c |1\rangle - \Omega_p |2\rangle$$

Destructive interference,
cancellation of absorption

— **EIT**

Effective Magnetic Field

$$\mathbf{A} = -\hbar \frac{|\zeta|^2}{1 + |\zeta|^2} \nabla S, \quad \mathbf{B} = \hbar \frac{\nabla S \times \nabla |\zeta|^2}{(1 + |\zeta|^2)^2},$$
$$\phi = \frac{\hbar^2}{2M} \frac{(\nabla |\zeta|)^2 + |\zeta|^2 (\nabla S)^2}{(1 + |\zeta|^2)^2},$$

where

$$\zeta = \Omega_p / \Omega_c = |\zeta| e^{iS}.$$

- Light beams with relative OAM can introduce an effective magnetic field which acts on the electrically neutral atoms.
- The vector potential \mathbf{A} is determined by:
 - the gradient of phase difference between the probe and control beams,
 - the ratio between the intensities of the control and probe beams.

Effective Magnetic Field

$$\mathbf{A} = -\hbar \frac{|\zeta|^2}{1 + |\zeta|^2} \nabla S, \quad \mathbf{B} = \hbar \frac{\nabla S \times \nabla |\zeta|^2}{(1 + |\zeta|^2)^2},$$
$$\phi = \frac{\hbar^2}{2M} \frac{(\nabla |\zeta|)^2 + |\zeta|^2 (\nabla S)^2}{(1 + |\zeta|^2)^2},$$

where

$$\zeta = \Omega_p / \Omega_c = |\zeta| e^{iS}.$$

- Light beams with relative OAM can introduce an effective magnetic field which acts on the electrically neutral atoms.
- The vector potential \mathbf{A} is determined by:
 - the gradient of phase difference between the probe and control beams,
 - the ratio between the intensities of the control and probe beams.

Effective Magnetic Field

$$\mathbf{A} = -\hbar \frac{|\zeta|^2}{1 + |\zeta|^2} \nabla S, \quad \mathbf{B} = \hbar \frac{\nabla S \times \nabla |\zeta|^2}{(1 + |\zeta|^2)^2},$$
$$\phi = \frac{\hbar^2}{2M} \frac{(\nabla |\zeta|)^2 + |\zeta|^2 (\nabla S)^2}{(1 + |\zeta|^2)^2},$$

where

$$\zeta = \Omega_p / \Omega_c = |\zeta| e^{iS}.$$

- Light beams with relative OAM can introduce an effective magnetic field which acts on the electrically neutral atoms.
- The vector potential \mathbf{A} is determined by:
 - the gradient of phase difference between the probe and control beams,
 - the ratio between the intensities of the control and probe beams.

Effective Magnetic Field


$$\mathbf{A} = -\hbar \frac{|\zeta|^2}{1 + |\zeta|^2} \nabla S, \quad \mathbf{B} = \hbar \frac{\nabla S \times \nabla |\zeta|^2}{(1 + |\zeta|^2)^2},$$
$$\phi = \frac{\hbar^2}{2M} \frac{(\nabla |\zeta|)^2 + |\zeta|^2 (\nabla S)^2}{(1 + |\zeta|^2)^2},$$

where

$$\zeta = \Omega_p / \Omega_c = |\zeta| e^{iS}.$$

- Light beams with relative OAM can introduce an effective magnetic field which acts on the electrically neutral atoms.
- The vector potential \mathbf{A} is determined by:
 - the gradient of phase difference between the probe and control beams,
 - the ratio between the intensities of the control and probe beams.

Light beams with OAM: Light Vortices



Light vortex

Light vortex — light beam with phase


$$e^{ikz+il\varphi},$$

where φ is azimuthal angle, l — winding number.

Light vortices have **orbital angular momentum** (OAM) along the propagation axis $M_z = \hbar l$.

- G. Juzeliūnas and P. Öhberg, Phys. Rev. Lett. **93**, 033602 (2004).
- G. Juzeliūnas, P. Öhberg, J. Ruseckas, and A. Klein, Phys. Rev. A **71**, 053614 (2005).


Counterpropagating Light Beams



The relative phase $S = (k_p + k_c)y$


J. Ruseckas, G. Juzeliūnas, P. Öhberg, and M. Fleischhauer, Phys. Rev. A **73** 025602 (2006).

Counterpropagating Gaussian Beams



Effective magnetic field B_{eff} and effective trapping potential $V_{\text{eff}} = V + \phi$ produced by counter-propagating Gaussian beams.

Other configurations



K. J. Günter, M. Cheneau, T. Yefsah, S. P. Rath, and J. Dalibard,
Practical scheme for a light-induced gauge field in an atomic Bose gas,
Phys. Rev. A **79**, 011604(R) (2009).

Effective magnetic field induced by position-dependent detuning

Alternative method

Effective gauge potentials also can be created using position-dependent **detuning**.

- The Hamiltonian for the electronic degrees of freedom $\hat{H}_0(\mathbf{r})$ includes position-dependent detuning $\delta(\mathbf{r})$.
- Using adiabatic approximation the same general expressions for the geometric potentials apply.

Effective magnetic field induced by position-dependent detuning

Alternative method

Effective gauge potentials also can be created using position-dependent **detuning**.

- The Hamiltonian for the electronic degrees of freedom $\hat{H}_0(\mathbf{r})$ includes position-dependent detuning $\delta(\mathbf{r})$.
- Using adiabatic approximation the same general expressions for the geometric potentials apply.

Effective magnetic field induced by position-dependent detuning


Alternative method

Effective gauge potentials also can be created using position-dependent **detuning**.


- The Hamiltonian for the electronic degrees of freedom $\hat{H}_0(\mathbf{r})$ includes position-dependent detuning $\delta(\mathbf{r})$.
- Using adiabatic approximation the same general expressions for the geometric potentials apply.

Experimental realization

(a) Experimental layout



(b) Level diagram




- Counterpropagating σ_+ and π laser beams
- Atom in a real magnetic field ($F=1$)
- Raman coupling between the ground states $m_F = \pm 1$ and $m_F = 0$.


Y.-J. Lin, R. L. Compton, A. R. Perry, W. D. Phillips, J. V. Porto, and I. B. Spielman,
Phys. Rev. Lett. **102**, 130401 (2009).

Experimental realization

(a) Experimental layout



(b) Level diagram




- Counterpropagating σ_+ and π laser beams
- Atom in a real magnetic field ($F=1$)
- Raman coupling between the ground states $m_F = \pm 1$ and $m_F = 0$.


Y.-J. Lin, R. L. Compton, A. R. Perry, W. D. Phillips, J. V. Porto, and I. B. Spielman, Phys. Rev. Lett. **102**, 130401 (2009).

Experimental realization

(a) Experimental layout



(b) Level diagram




- Counterpropagating σ_+ and π laser beams
- Atom in a real magnetic field ($F=1$)
- Raman coupling between the ground states $m_F = \pm 1$ and $m_F = 0$.


Y.-J. Lin, R. L. Compton, A. R. Perry, W. D. Phillips, J. V. Porto, and I. B. Spielman,
Phys. Rev. Lett. **102**, 130401 (2009).

Experimental realization

(a) Experimental layout



(b) Level diagram



- Counterpropagating σ_+ and π laser beams
- Atom in a real magnetic field ($F=1$)
- Raman coupling between the ground states $m_F = \pm 1$ and $m_F = 0$.

Y.-J. Lin, R. L. Compton, A. R. Perry, W. D. Phillips, J. V. Porto, and I. B. Spielman, Phys. Rev. Lett. **102**, 130401 (2009).

Experimental realization

An **alternative** description to the one used by Lin *et al.*

The Hamiltonian for the electronic degrees of freedom

$$H_0 = \hbar \begin{pmatrix} -\delta & \Omega_R^* & 0 \\ \Omega_R & 0 & \Omega_R^* \\ 0 & \Omega_R & \delta \end{pmatrix}$$

with two-photon coupling $\Omega_R = |\Omega| e^{ik_d x}$.

Atom stays in the lowest-energy eigenstate

$$|\chi_{-}\rangle = e^{-ik_d x} \cos^2(\theta/2) |-1\rangle - 1/\sqrt{2} \sin \theta |0\rangle + e^{ik_d x} \sin^2(\theta/2) |1\rangle$$

where $\theta \equiv \arctan(\sqrt{2}|\Omega|/\delta)$.

The effective vector potential

$$\mathbf{A} = \hbar k_d \cos \theta \mathbf{e}_x \approx \hbar k_d \delta / (\sqrt{2}|\Omega|) \mathbf{e}_x$$

Experimental realization

An **alternative** description to the one used by Lin *et al.*.
The Hamiltonian for the electronic degrees of freedom

$$H_0 = \hbar \begin{pmatrix} -\delta & \Omega_R^* & 0 \\ \Omega_R & 0 & \Omega_R^* \\ 0 & \Omega_R & \delta \end{pmatrix}$$

with two-photon coupling $\Omega_R = |\Omega| e^{ik_d x}$.

Atom stays in the lowest-energy eigenstate

$$|\chi_{-}\rangle = e^{-ik_d x} \cos^2(\theta/2) |-1\rangle - 1/\sqrt{2} \sin \theta |0\rangle + e^{ik_d x} \sin^2(\theta/2) |1\rangle$$

where $\theta \equiv \arctan(\sqrt{2}|\Omega|/\delta)$.

The effective vector potential

$$\mathbf{A} = \hbar k_d \cos \theta \mathbf{e}_x \approx \hbar k_d \delta / (\sqrt{2}|\Omega|) \mathbf{e}_x$$

Experimental realization

An **alternative** description to the one used by Lin *et al.*.
The Hamiltonian for the electronic degrees of freedom

$$H_0 = \hbar \begin{pmatrix} -\delta & \Omega_R^* & 0 \\ \Omega_R & 0 & \Omega_R^* \\ 0 & \Omega_R & \delta \end{pmatrix}$$

with two-photon coupling $\Omega_R = |\Omega| e^{ik_d x}$.

Atom stays in the lowest-energy eigenstate

$$|\chi_{-}\rangle = e^{-ik_d x} \cos^2(\theta/2) | -1 \rangle - 1/\sqrt{2} \sin \theta |0\rangle + e^{ik_d x} \sin^2(\theta/2) |1\rangle$$

where $\theta \equiv \arctan(\sqrt{2}|\Omega|/\delta)$.

The effective vector potential

$$\mathbf{A} = \hbar k_d \cos \theta \mathbf{e}_x \approx \hbar k_d \delta / (\sqrt{2}|\Omega|) \mathbf{e}_x$$

Experimental realization

An **alternative** description to the one used by Lin *et al.*
The Hamiltonian for the electronic degrees of freedom

$$H_0 = \hbar \begin{pmatrix} -\delta & \Omega_R^* & 0 \\ \Omega_R & 0 & \Omega_R^* \\ 0 & \Omega_R & \delta \end{pmatrix}$$

with two-photon coupling $\Omega_R = |\Omega| e^{ik_d x}$.

Atom stays in the lowest-energy eigenstate

$$|\chi_{-}\rangle = e^{-ik_d x} \cos^2(\theta/2) |-1\rangle - 1/\sqrt{2} \sin \theta |0\rangle + e^{ik_d x} \sin^2(\theta/2) |1\rangle$$


where $\theta \equiv \arctan(\sqrt{2}|\Omega|/\delta)$.

The effective vector potential


$$\mathbf{A} = \hbar k_d \cos \theta \mathbf{e}_x \approx \hbar k_d \delta / (\sqrt{2}|\Omega|) \mathbf{e}_x$$

Experimental realization


a Geometry



b Level diagram



Dressed state, $\hbar\Omega_R = 8.20E_L$



Y.-J. Lin, R. L. Compton,
K. Jiménez-García, J. V. Porto and
I. B. Spielman, *Nature*, **462**, 628 (2009).


Non-Abelian gauge potentials

- Adiabatic motion of many-level cold atoms in the laser fields varying in space creates effective **non-Abelian** gauge fields.
- It is possible to simulate motion of the particles in the non-Abelian fields using cold atomic gases.


Non-Abelian gauge potentials

- Adiabatic motion of many-level cold atoms in the laser fields varying in space creates effective **non-Abelian** gauge fields.
- It is possible to simulate motion of the particles in the non-Abelian fields using cold atomic gases.

Tripod Coupling Scheme




Tripod Coupling Scheme



- Two degenerate dark states
- Non-Abelian gauge potentials

- R. G. Unanyan, M. Fleischhauer, B. E. Shore, and K. Bergmann, *Opt. Commun.* **155**, 144 (1998).
- J. Ruseckas, G. Juzeliūnas, P. Öhberg, and M. Fleischhauer *Phys. Rev. Lett.* **95**, 010404 (2005)


Tripod Coupling Scheme



- Two degenerate dark states
- Non-Abelian gauge potentials

- R. G. Unanyan, M. Fleischhauer, B. E. Shore, and K. Bergmann, *Opt. Commun.* **155**, 144 (1998).
- J. Ruseckas, G. Juzeliūnas, P. Öhberg, and M. Fleischhauer *Phys. Rev. Lett.* **95**, 010404 (2005)

Tripod Coupling Scheme



- **Two** degenerate dark states
- **Non-Abelian** gauge potentials

- R. G. Unanyan, M. Fleischhauer, B. E. Shore, and K. Bergmann, *Opt. Commun.* **155**, 144 (1998).
- J. Ruseckas, G. Juzeliūnas, P. Öhberg, and M. Fleischhauer *Phys. Rev. Lett.* **95**, 010404 (2005)

Magnetic Monopole

- Laser fields:

$$\Omega_{1,2} = \Omega_0 \frac{\rho}{R} e^{i(kz \mp \varphi)}, \quad \Omega_3 = \Omega_0 \frac{z}{R} e^{ik'x}.$$

- The effective magnetic field

$$\mathbf{B} = \frac{\hbar}{r^2} \mathbf{e}_r \begin{pmatrix} 0 & 1 \\ 1 & 0 \end{pmatrix} + \dots$$

- J. Ruseckas, G. Juzeliūnas, P. Öhberg, and M. Fleischhauer, Phys. Rev. Lett. **95**, 010404 (2005).
- V. Pietila, M. Mottonen, Phys. Rev. Lett. **102**, 080403 (2009).
- V. Pietila, M. Mottonen, Phys. Rev. Lett. **103**, 030401 (2009).

Magnetic Monopole

- Laser fields:


$$\Omega_{1,2} = \Omega_0 \frac{\rho}{R} e^{i(kz \mp \varphi)}, \quad \Omega_3 = \Omega_0 \frac{z}{R} e^{ik'x}.$$

- The effective magnetic field

$$\mathbf{B} = \frac{\hbar}{r^2} \mathbf{e}_r \begin{pmatrix} 0 & 1 \\ 1 & 0 \end{pmatrix} + \dots$$

- J. Ruseckas, G. Juzeliūnas, P. Öhberg, and M. Fleischhauer, Phys. Rev. Lett. **95**, 010404 (2005).
- V. Pietila, M. Mottonen, Phys. Rev. Lett. **102**, 080403 (2009).
- V. Pietila, M. Mottonen, Phys. Rev. Lett. **103**, 030401 (2009).

Ultrarelativistic Dirac fermions



$$\Omega_1 = \Omega \sin \theta e^{-i\kappa x} / \sqrt{2}, \quad \Omega_2 = \Omega \sin \theta e^{i\kappa x} / \sqrt{2}, \quad \Omega_3 = \Omega \cos \theta e^{-i\kappa y}$$

where

$$\theta = \theta_0, \quad \cos \theta_0 = \sqrt{2} - 1$$

Ultrarelativistic Dirac fermions

The Hamiltonian

$$H_{\mathbf{k}} = \frac{\hbar^2}{2m} (\mathbf{k} + \kappa' \sigma_{\perp})^2 + V_1$$

with

$$\kappa' = \kappa \cos \theta_0, \quad \sigma_{\perp} = \mathbf{e}_x \sigma_x + \mathbf{e}_y \sigma_y$$

For small wave vectors $k \ll \kappa'$, the atomic Hamiltonian reduces to the Hamiltonian for the 2D relativistic motion of a two-component massless particle of the Dirac type known also as the **Weyl equation**

$$H_{\mathbf{k}} = \hbar v_0 \mathbf{k} \cdot \sigma_{\perp} + V_1 + mv_0^2$$

where the velocity $v_0 = \hbar \kappa' / m$ corresponds to the velocity of light. For cold atoms this velocity is of the order 1 cm/s.

Ultrarelativistic Dirac fermions

The Hamiltonian

$$H_{\mathbf{k}} = \frac{\hbar^2}{2m} (\mathbf{k} + \kappa' \sigma_{\perp})^2 + V_1$$

with


$$\kappa' = \kappa \cos \theta_0, \quad \sigma_{\perp} = \mathbf{e}_x \sigma_x + \mathbf{e}_y \sigma_y$$

For small wave vectors $k \ll \kappa'$, the atomic Hamiltonian reduces to the Hamiltonian for the 2D relativistic motion of a two-component massless particle of the Dirac type known also as the **Weyl equation**

$$H_{\mathbf{k}} = \hbar v_0 \mathbf{k} \cdot \sigma_{\perp} + V_1 + mv_0^2$$

where the velocity $v_0 = \hbar \kappa' / m$ corresponds to the velocity of light. For cold atoms this velocity is of the order 1 cm/s.

Ultrarelativistic Dirac fermions



The Hamiltonian for small momenta with an additional scalar potential:

$$H = v_0 \sigma_{\perp} \cdot \mathbf{p} + V \sigma_z$$

The velocity operator

$$\mathbf{v} \equiv \dot{\mathbf{r}} = \frac{1}{i\hbar} [\mathbf{r}, H] = v_0 \sigma_{\perp}$$

The eigenfunctions of the Hamiltonian do not have a definite velocity.

Consequence: oscillations in the movement of the wave packet.

- J. Y. Vaishnav and C. W. Clark, Phys. Rev. Lett. **100**, 153002 (2008).
- M. Merkl, F. E. Zimmer, G. Juzeliūnas, and P. Öhberg, Europhys. Lett. **83**, 54002 (2008).

The Hamiltonian for small momenta with an additional scalar potential:

$$H = v_0 \sigma_{\perp} \cdot \mathbf{p} + V \sigma_z$$

The velocity operator

$$\mathbf{v} \equiv \dot{\mathbf{r}} = \frac{1}{i\hbar} [\mathbf{r}, H] = v_0 \sigma_{\perp}$$

The eigenfunctions of the Hamiltonian do not have a definite velocity.
Consequence: oscillations in the movement of the wave packet.

- J. Y. Vaishnav and C. W. Clark, Phys. Rev. Lett. **100**, 153002 (2008).
- M. Merkl, F. E. Zimmer, G. Juzeliūnas, and P. Öhberg, Europhys. Lett. **83**, 54002 (2008).

The Hamiltonian for small momenta with an additional scalar potential:

$$H = v_0 \sigma_{\perp} \cdot \mathbf{p} + V \sigma_z$$

The velocity operator

$$\mathbf{v} \equiv \dot{\mathbf{r}} = \frac{1}{i\hbar} [\mathbf{r}, H] = v_0 \sigma_{\perp}$$

The eigenfunctions of the Hamiltonian do not have a definite velocity.

Consequence: oscillations in the movement of the wave packet.

- J. Y. Vaishnav and C. W. Clark, Phys. Rev. Lett. **100**, 153002 (2008).
- M. Merkl, F. E. Zimmer, G. Juzeliūnas, and P. Öhberg, Europhys. Lett. **83**, 54002 (2008).

The Hamiltonian for small momenta with an additional scalar potential:

$$H = v_0 \sigma_{\perp} \cdot \mathbf{p} + V \sigma_z$$

The velocity operator


$$\mathbf{v} \equiv \dot{\mathbf{r}} = \frac{1}{i\hbar} [\mathbf{r}, H] = v_0 \sigma_{\perp}$$

The eigenfunctions of the Hamiltonian do not have a definite velocity.


Consequence: oscillations in the movement of the wave packet.

- J. Y. Vaishnav and C. W. Clark, Phys. Rev. Lett. **100**, 153002 (2008).
- M. Merkl, F. E. Zimmer, G. Juzeliūnas, and P. Öhberg, Europhys. Lett. **83**, 54002 (2008).

Negative reflection




Negative reflection




Reflection probabilities.

Negative reflection




G. Juzeliūnas, J. Ruseckas, A. Jacob, L. Santos, and P. Öhberg, Phys. Rev. Lett. **100**, 200405 (2008).

Spin field effect transistor with ultracold atoms



J. Y. Vaishnav, J. Ruseckas, C. W. Clark, and G. Juzeliūnas, Phys. Rev. Lett. **101**, 265302 (2008).


Tetrapod scheme with counter-propagating beams




Lambda-type scheme, no Raman coupling to the $F = 2$ levels

G. Juzeliūnas, J. Ruseckas, and J. Dalibard, Phys. Rev. A **81**, 053403 (2010).

Tetrapod scheme with counter-propagating beams




Lambda-type scheme, no Raman coupling to the $F=2$ levels






Raman coupling to the $F=2$ levels

G. Juzeliūnas, J. Ruseckas, and J. Dalibard, Phys. Rev. A **81**, 053403 (2010).

Tetrapod scheme with counter-propagating beams



Tetrapod scheme with counter-propagating beams



Tetrapod scheme with counter-propagating beams

Spin-1 **Rashba-type** Hamiltonian

$$\hat{H} = \frac{1}{2m}(\hat{\mathbf{p}} + \hbar\kappa\mathbf{J}_{\perp})^2 + V$$


where \mathbf{J}_{\perp} is the projection of spin-1 operator onto the xy plane.

Tetrapod scheme with counter-propagating beams

Spin-1 **Rashba-type** Hamiltonian


$$\hat{H} = \frac{1}{2m}(\hat{\mathbf{p}} + \hbar\kappa\mathbf{J}_{\perp})^2 + V$$

where \mathbf{J}_{\perp} is the projection of spin-1 operator onto the xy plane.



Dispersion

Comparison of transmission probabilities for spin-1/2 and spin-1 systems



Summary

- Light beams with relative orbital angular momentum can introduce Abelian and non-Abelian effective gauge potentials acting on the electrically neutral atoms.
- Non-Abelian fields can be formed for cold atoms using the plane-wave setups. This was not possible for the Abelian fields.
- Atomic motion in non-Abelian fields exhibits a number of non-trivial features, such as their quasirelativistic behavior or the negative refraction and reflection.
- The plane wave setups can lead to the spin $1/2$ or the spin 1 Rashba-type Hamiltonian for cold atoms.

Summary

- Light beams with relative orbital angular momentum can introduce Abelian and non-Abelian effective gauge potentials acting on the electrically neutral atoms.
- Non-Abelian fields can be formed for cold atoms using the plane-wave setups. This was not possible for the Abelian fields.
- Atomic motion in non-Abelian fields exhibits a number of non-trivial features, such as their quasirelativistic behavior or the negative refraction and reflection.
- The plane wave setups can lead to the spin $1/2$ or the spin 1 Rashba-type Hamiltonian for cold atoms.

Summary

- Light beams with relative orbital angular momentum can introduce Abelian and non-Abelian effective gauge potentials acting on the electrically neutral atoms.
- Non-Abelian fields can be formed for cold atoms using the plane-wave setups. This was not possible for the Abelian fields.
- Atomic motion in non-Abelian fields exhibits a number of non-trivial features, such as their quasirelativistic behavior or the negative refraction and reflection.
- The plane wave setups can lead to the spin $1/2$ or the spin 1 Rashba-type Hamiltonian for cold atoms.

- Light beams with relative orbital angular momentum can introduce Abelian and non-Abelian effective gauge potentials acting on the electrically neutral atoms.
- Non-Abelian fields can be formed for cold atoms using the plane-wave setups. This was not possible for the Abelian fields.
- Atomic motion in non-Abelian fields exhibits a number of non-trivial features, such as their quasirelativistic behavior or the negative refraction and reflection.
- The plane wave setups can lead to the spin $1/2$ or the spin 1 Rashba-type Hamiltonian for cold atoms.

Thank you!